



## Section 1.1 – Real Numbers



### لاحظ

Rational Numbers ( $\mathbb{Q}$ ) can be expressed in three formats:

- Fraction

$$\frac{1}{2}$$

$$-\frac{3}{7}$$

$$46 = \frac{46}{1}$$

- Finite decimal numbers

$$0.17$$

$$0.5$$

- Repeating decimal representation

$$0.\overline{6} = 0.66666 \dots$$

$$0.3\overline{17} = 0.3171717 \dots$$

## Example 1

Express the decimal  $0.2\overline{8}$  as a fraction.

## Solution

$$x = 0.2\overline{8} = 0.28888\ldots$$

$$100x = 28.8888\ldots$$

$$10x = 2.8888\ldots$$

$$\begin{array}{r} 90x = 26 \\ \hline \end{array}$$

$$\boxed{x = \frac{26}{90}}$$

- Properties of real numbers خصائص الأعداد الحقيقية

## Commutative

$$a + b = b + a$$

$$3 + 4 = 4 + 3$$

$$ab = ba$$

$$5 \cdot 6 = 6 \cdot 5$$

## Associative

$$(a + b) + c = a + (b + c)$$

$$(2 + 3) + 7 = 2 + (3 + 7)$$

$$(ab)c = a(bc)$$

$$(4 \cdot 5) \cdot 3 = 4 \cdot (5 \cdot 3)$$

## Distributive

$$a(b + c) = ab + ac$$

$$2 \cdot (6 + 4) = 2 \cdot 6 + 2 \cdot 4$$

$$(b + c)a = ab + ac$$

$$(6 + 4) \cdot 2 = 2 \cdot 6 + 2 \cdot 4$$

### Example 2

Use properties of real numbers to write the expression without parentheses:

(a)  $3(x + y)$

(b)  $(3a)(b + c - 2d)$

**Solution**

$$(a) \quad 3(\overset{\curvearrowright}{\underset{\curvearrowright}{x+y}}) = 3x + 3y$$

$$(b) \quad (3a)(\overset{\curvearrowright}{\underset{\curvearrowright}{b+c-2d}}) = 3ab + 3ac - 6ad$$

- The number 0 is called the **additive identity** المحاييد الجمعي because  $a + 0 = a$  for any real number  $a$ .

- **Properties of negatives**

1.  $(-1)a = -a$

2.  $-(-a) = a$

3.  $(-a)b = a(-b) = -(ab)$

4.  $(-a)(-b) = ab$

5.  $-(a + b) = -a - b$

6.  $-(a - b) = b - a$

### Example 3

Use properties of negatives to write the expression  $-(x + y - z)$  without parentheses.

**Solution**

$$-x - y + z$$

- The number 1 is called the **multiplicative identity** المحاييد الضربي because  $a \cdot 1 = a$  for any real number  $a$ .

- Division القسمة is the inverse of multiplication, hence:

$$a \div b = a \cdot \frac{1}{b}$$

$$= \frac{a}{b}$$

Numerator البسط

Denominator المقام

### - Properties of fractions

$$1. \frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

$$\frac{2}{3} \cdot \frac{5}{7} = \frac{2 \cdot 5}{3 \cdot 7} = \frac{10}{21}$$

$$2. \frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}$$

$$\frac{2}{3} \div \frac{5}{7} = \frac{2}{3} \cdot \frac{7}{5} = \frac{14}{15}$$

$$3. \frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

$$\frac{2}{5} + \frac{7}{5} = \frac{2+7}{5} = \frac{9}{5}$$

$$4. \frac{a}{b} + \frac{c}{d} = \frac{ad+b}{bd}$$

$$\frac{2}{5} + \frac{3}{7} = \frac{2 \cdot 7 + 3 \cdot 5}{5 \cdot 7} = \frac{14+15}{35} = \frac{29}{35}$$

$$5. \frac{ac}{bc} = \frac{a}{b}$$

$$\frac{2 \cdot 5}{3 \cdot 5} = \frac{2}{3}$$

$$6. \text{ If } \frac{a}{b} = \frac{c}{d}, \text{ then } ad = bc$$

$$\frac{2}{3} = \frac{6}{9}, \text{ so } 2 \cdot 9 = 3 \cdot 6$$

## Example 4

Perform the indicated operations.

(a)  $\frac{3}{10} + \frac{4}{15}$

(b)  $\frac{2}{3}(6 - \frac{3}{2})$

(c)  $\frac{\frac{2}{5} + \frac{1}{2}}{\frac{1}{10} + \frac{3}{15}}$

**Solution**

$$\begin{aligned} \text{(a)} \quad \frac{3}{10} + \frac{4}{15} \\ &= \frac{3 \times 15 + 4 \times 10}{10 \times 15} = \frac{45 + 40}{150} \\ &= \frac{85}{150} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \frac{2}{3} \left( 6 - \frac{3}{2} \right) &= \frac{2}{3} \cdot 6 - \frac{2}{3} \cdot \frac{3}{2} \\ &= \frac{12}{3} - \frac{6}{6} = 4 - 1 = 3 \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \frac{\frac{2}{5} + \frac{1}{2}}{\frac{1}{10} + \frac{3}{15}} &= \left( \frac{2}{5} + \frac{1}{2} \right) \div \left( \frac{1}{10} + \frac{3}{15} \right) \\ &= \left( \frac{2 \times 2 + 1 \times 5}{5 \times 2} \right) \div \left( \frac{1 \times 15 + 3 \times 10}{10 \times 15} \right) \\ &= \left( \frac{9}{10} \right) \div \left( \frac{45}{150} \right) \\ &= \frac{9}{10} \times \frac{150}{45} = \frac{1350}{450} = 3 \end{aligned}$$

- If  $a$  is a real number, then the **absolute value** مطلق of  $a$  is

$$|a| = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases}$$

- **Properties of absolute values**

$$1. |a| \geq 0 \qquad |-3| = 3 \geq 0$$

$$2. |a| = |-a| \qquad |5| = |-5|$$

$$3. |ab| = |a||b| \qquad |-3 \cdot 5| = |-3||5|$$

$$4. \left| \frac{a}{b} \right| = \frac{|a|}{|b|} \qquad \left| \frac{12}{-3} \right| = \frac{|12|}{|-3|}$$

$$5. |a + b| \leq |a| + |b| \qquad |-3 + 5| \leq |-3| + |5|$$

### Example 5

Evaluate each expression.

$$(a) ||-6| - |-4||$$

$$(b) \left| \frac{7-12}{12-7} \right|$$

**Solution**

$$(a) |6 - 4| = |2| = 2$$

$$(b) \left| \frac{-5}{5} \right| = |-1| = 1$$

- The distance between any two points  $a$  and  $b$  on the real line is

$$d(a, b) = |b - a| = |a - b|$$

### Example 6

Find the distance between  $-8$  and  $2$

**Solution**

$$D = |-8 - 2| = |-10| = 10$$

- A set المجموعة is a collection of objects called elements العناصر

$$A = \{1,2,3,4,5,6\}$$

لاحظ الأقواس المستخدمة { }

$$A = \{x | x \text{ is an integer and } 0 < x < 7\}$$

طريقة أخرى لكتابة

المجموعات تسمى

**set-builder notation**

- If  $A$  and  $B$  are sets, then their **union** اتحاد  $A \cup B$  is the set that consists of all the elements that are in  $A$  and  $B$

Example:

$$A = \{1,2,3,4,5\} \text{ and } B = \{4,5,6,7\}$$

$$A \cup B = \{1,2,3,4,5,6,7\}$$

- The **intersection** التقاطع of  $A$  and  $B$  is the set  $A \cap B$  consisting of all elements that are in both  $A$  and  $B$ .

Example:

$$A = \{1,2,3,4,5\} \text{ and } B = \{4,5,6,7\}$$

$$A \cap B = \{4,5\}$$

- إذا كانت المجموعة ليس بها عناصر تسمى مجموعة خالية **empty set** و يرمز لها بالرمز  $\emptyset$  (Phi)

Example:

$$A = \{1,2,3\} \text{ and } B = \{4,5,6,7\}$$

$$A \cap B = \emptyset$$

- **Intervals** الفترات are sets with an infinite number of elements and are defined by start and end values.

Open interval



$$(a, b) = \{x | a < x < b\}$$

Closed interval



$$[a, b] = \{x | a \leq x \leq b\}$$

Notation	Set description	Graph
$(a, b)$	$\{x   a < x < b\}$	
$[a, b]$	$\{x   a \leq x \leq b\}$	
$[a, b)$	$\{x   a \leq x < b\}$	
$(a, b]$	$\{x   a < x \leq b\}$	
$(a, \infty)$	$\{x   a < x\}$	
$[a, \infty)$	$\{x   a \leq x\}$	
$(-\infty, b)$	$\{x   x < b\}$	
$(-\infty, b]$	$\{x   x \leq b\}$	
$(-\infty, \infty)$	$\mathbb{R}$ (set of all real numbers)	


### Example 7

Graph each set.

(a)  $(1, 3) \cap [2, 7]$

(b)  $(1, 3) \cup [2, 7]$

### Solution





**Problems**

- Perform the indicated operations.

$$(a) \frac{\frac{2}{2}}{\frac{3}{3}} - \frac{\frac{2}{3}}{2}$$

$$\begin{aligned} & \left( 2 \div \frac{2}{3} \right) - \left( \frac{2}{3} \div 2 \right) \\ &= \left( \cancel{2} \cdot \frac{3}{\cancel{2}} \right) - \left( \frac{\cancel{2}}{3} \cdot \frac{1}{\cancel{2}} \right) \\ &= 3 - \frac{1}{3} \\ &= \frac{3 \cdot 3 - 1}{3} = \frac{8}{3} \end{aligned}$$

$$(b) 1 + \frac{5}{8} - \frac{1}{6}$$

$$\begin{aligned} &= \frac{8+5}{8} - \frac{1}{6} = \frac{13}{8} - \frac{1}{6} \\ &= \frac{13 \cdot 6 - 1 \cdot 8}{8 \cdot 6} = \frac{78 - 8}{48} \\ &= \frac{70}{48} = \frac{35}{24} \end{aligned}$$

$$(c) \left( 3 + \frac{1}{4} \right) \left( 1 - \frac{4}{5} \right)$$

$$\begin{aligned} & \left( \frac{3 \cdot 4 + 1}{4} \right) \left( \frac{1 \cdot 5 - 4}{5} \right) \\ &= \frac{12+1}{4} \cdot \frac{5-4}{5} \\ &= \frac{13}{4} \cdot \frac{1}{5} = \frac{13}{20} \end{aligned}$$

- Find the indicated set if

$$A = \{1, 2, 3, 4, 5, 6, 7\}, B = \{2, 4, 6, 8\}, \text{ and } C = \{7, 8, 9, 10\}$$

(a)  $A \cup B \cup C$

$$= \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

(b)  $A \cap B \cap C = \emptyset$


empty set

- Find the indicated set if


$$A = \{x | x \geq -2\}, B = \{x | x < 4\}, \text{ and } C = \{x | -1 < x \leq 5\}$$

(a)  $B \cup C = \{x | x \leq 5\}$

يمكن ترتيب لتخيل الحل




(b)  $A \cap C = \{x | -1 < x \leq 5\}$



- Express the inequality in interval notation, and then graph the corresponding interval.

(a)  $x \leq 1$

$$x = (-\infty, 1]$$



(b)  $-2 < x \leq 1$

$$x = (-2, 1]$$



- Find the distance between the given numbers.

(a)  $-3$  and  $21$

$$\begin{aligned} d &= |-3 - 21| \\ &= |-24| = 24 \end{aligned}$$

(b)  $\frac{11}{8}$  and  $-\frac{3}{10}$

$$\begin{aligned} d &= \left| \frac{11}{8} - \left(-\frac{3}{10}\right) \right| = \left| \frac{11}{8} + \frac{3}{10} \right| \\ &= \left| \frac{11 \cdot 10 + 3 \cdot 8}{80} \right| = \left| \frac{110 + 24}{80} \right| = \frac{134}{80} \end{aligned}$$

- Express each repeating decimal as a fraction

(a)  $0.2\overline{8} = 0.28888 \dots$

$$\begin{aligned} 100x &= 28.88 \\ 10x &= 2.888 \\ \hline 90x &= 26 \\ \boxed{x} &= \frac{26}{90} \end{aligned}$$

(b)  $5.\overline{23} = 5.232323 \dots$

$$\begin{aligned} 100x &= 523.23 \\ x &= 5.23 \\ \hline 99x &= 518 \\ \boxed{x} &= \frac{518}{99} \end{aligned}$$

(c)  $2.1\overline{35} = 2.1353535 \dots$

$$\begin{aligned} 1000x &= 2135.35 \\ 10x &= 21.3535 \\ \hline 990x &= 2114 \\ \boxed{x} &= \frac{2114}{990} \end{aligned}$$